


Bedrock Geology of the Portland 1:100,000 Quadrangle, Maine and New Hampshire



Topographic base from U.S. Geological Survey Portland, Maine - New Hampshire 1:100,000-scale metric topographic-bathymetric map

Miles (mi) 0 1 2 3 4 5 6 7 8 9 10 11 12

Map Scale 1:100,000

Kilometers (km) 0 2 4 6 8 10 12 14 16 18 20

Contour interval 10 meters. National geodetic vertical datum of 1929. Elevations shown to nearest meter.

INTRUSIVE AND RELATED VOLCANIC ROCKS

MESOZOIC

Randall Mountain Stock

- Mt** Fragmental trachyte porphyry.
- Mbs** Equigranular to porphyritic biotite-hornblende alkali feldspar syenite.

Burnt Meadow Mountains Igneous Complex

- Mbv** Volcanic breccia and feldspar porphyry. May include shallow intrusive rocks.
- Mbs** Alkali feldspar syenite. Brown to gray.
- Mbsq** Quartz syenite. Pinkish-tan.

Acton Stock

- Ma** Dark gray, porphyritic and fragmental andesite.
- Md** Medium-grained pyroxene-quartz diorite.

Rattlesnake Mountain Igneous Complex

- Mre** Nepheline syenite.
- Mrd** Nepheline-bearing syenite.
- Mrc** Fine-grained syenite.
- Mrb** Ferrobastingsite syenite.
- Mra** Biotite-ferrobastingsite syenite.

Abbott Mountain Stock

- Mas** Light colored syenite.
- Mal** Fayalite-pyroxene syenite.
- Maq** Fayalite-pyroxene-quartz syenite.

Other Rocks of the White Mountain Magma Series

- Ms** Fayalite-pyroxene syenite (at Symmes Pond, Boston Hills).
- Mbg** Medium-grained biotite-hornblende granite (at Picket Mountain).
- Mbg** Pink, biotite granite. Locally with fine-grained or porphyritic texture (equivalent to the Conway Granite, New Hampshire).

CARBONIFEROUS

Sebago Pluton

- Csg** Muscovite-biotite granite.
- Csb** Biotite granite, non-foliated.
- Csf** Biotite granite, foliated.
- Csgg** Muscovite-garnet granite.
- Csm** Mixed muscovite-garnet and muscovite-biotite granites.
- Csx** Migmatite muscovite-biotite granite.
- Csgr** Ribbeckite-bearing granite (near Rattlesnake Mountain).

CARBONIFEROUS(?)

Lyman Pluton

- Clg** Biotite-muscovite granite and pegmatite.

Other Plutonic Rocks

- Cg** Granites, mainly muscovite-bearing.
- Cp** Pegmatite, commonly with muscovite, garnet, and black tourmaline.

CARBONIFEROUS(?) OR DEVONIAN(?)

Saco Pluton

- CDsgb** Dark greenish-gray metamorphosed gabbro.
- CDsgd** Hornblende-biotite granodiorite.

Other Plutonic Rocks

- CDg** Granites, mainly muscovite-bearing.
- CDgb** Biotite granite.
- CDp** Pegmatite, commonly with muscovite, garnet, and black tourmaline.
- CDgp** Mixed granite and pegmatite.
- CDd** Dark greenish-gray diorite and gabbro.

DEVONIAN(?)

- Dgn** Gray gneiss: Foliated biotite-hornblende granodiorite.
- Dgbf** Foliated or lineated biotite granite.
- Dbgd** Gray biotite granodiorite.

STRATIFIED ROCKS

Shapleigh Group

- DSsq** Unnamed silvery gray, well bedded sillimanite-garnet-mica schist and quartzite (near Acton).

Sr

- Sr** Rindgemere Formation, undifferentiated: Reddish-brown to gray weathering feldspathic sillimanite-garnet-mica schist and migmatite are most common. Various other types of schist and granofels occur also.

Srlm

- Srlm** Rusty weathering schist.

Srlm

- Srlm** Libby Mountain member: Silvery gray, thin-bedded to medium-bedded sillimanite-garnet-mica schist and quartzite.

Srlm

- Srlm** Thick-bedded, white quartzite and subordinate quartz-mica schist.

Srlm

- Srlm** Thin-bedded sillimanite-garnet-mica schist and quartzite.

Srlm

- Srlm** Poorly to moderately well bedded, silvery gray, mica schist and quartzite (near North Alfred).

Srlm

- Srlm** Brown-weathering, feldspathic biotite schist.

Srlm

- Srlm** Muscovite-biotite-feldspar gneiss.

Srlm

- Srlm** Quartz-feldspar-biotite-garnet granofels.

Srlm

- Srlm** Interlayered calc-silicate granofels and granular biotite-quartz-feldspar schist. Calcic garnet (grossularite) and vesuvianite are present in some places.

Srlm

- Srlm** Quartz-feldspar-biotite-garnet granofels with interlayered calc-silicate granofels.

Srlm

- Srlm** Coarse-grained, migmatite muscovite-biotite-feldspar schist.

Srlm

- Srlm** Standish Formation, undifferentiated: Massive to well layered, gray to rusty weathering, sillimanite-garnet-mica schist interlayered with feldspathic granofels.

Srlm

- Srlm** Rusty weathering sillimanite-bearing schist, locally migmatite.

Srlm

- Srlm** Silvery-gray muscovite-rich schist.

Srlm

- Srlm** Thinly laminated calc-silicate and biotite-quartz-feldspar granofels and chlorite-biotite phyllite.

Central Maine Sequence

- SOmp** Unnamed migmatitic muscovite-bearing schist.

- SOmu** Undifferentiated migmatitic schist, gneiss, and granofels.

- SOms** Unnamed massive, brown-weathering, carbonate-rich schist and minor feldspathic granofels.

- SW** Windham Formation: Massive to thinly bedded biotite-muscovite-garnet-quartzite granofels with thin interbeds of greenish-gray calc-silicate granofels. Where migmatite, pegmatite layers commonly conform to bedding. May contain minor pelitic schist layers, but not generally in mappable amounts.

- SWl** Limestone member: Laminated to thinly layered impure marble, calc-silicate granofels and minor quartzite.

- SOhc** Hutchins Corner Formation: Flaggy, bluish to purplish-gray, biotite-quartz-plagioclase granofels with thin interbeds of greenish-gray calc-silicate granofels. Where migmatite, pegmatite layers commonly conform to bedding. May contain minor pelitic schist layers, but not generally in mappable amounts.

- SOhc1** Impure marble and calc-silicate rock.

- SOhc2** Pelitic schist, gray weathering.

- SOhc3** Rusty weathering schist.

- SOhc4** Chalky weathering, calcareous feldspathic gneiss and granofels. May be metamorphosed volcanic unit.

- SOhc5** Dipsoidal-plagioclase calc-silicate granofels.

- SOrc** Richmond Corner Formation: Garnetiferous quartz-plagioclase-biotite granular schist with minor amphibolite and garnet-quartz-magnetite granofels.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

- SOth** Toney Hill Formation: Extremely rusty weathering, sulfidic, graphitic schist.

Casco Bay Group

- OSpj** Spring Point to Jewell Formations, undifferentiated.

- Oj** Jewell Formation: Light gray to dark gray muscovite-biotite-garnet phyllite to schist, locally carbonaceous or rusty weathering. (Lithologically like the Scarborough Formation.)

- Ojf** Brownish-gray, chalky weathering, quartz-biotite phyllite.

- Osk** Spurvink Metalmistone: Fine-grained, metamorphosed limestone with thin interbeds of calcareous biotite-quartz phyllite. Contorted.

- Osc** Scarborough Formation: Light gray to dark gray muscovite-biotite-garnet phyllite to schist, locally carbonaceous or rusty weathering. (Lithologically like the Jewell Formation.)

- Osc1** Thin-bedded, fine-grained metamorphosed limestone and biotite phyllite.

- Osc2** Greenish-gray quartz-plagioclase-chlorite-biotite-garnet phyllite.

- Odi** Diamond Island Formation: Black, rusty weathering, quartz-muscovite-graphitic pyrite phyllite.

- Osp** Spring Point Formation: Greenish-gray plagioclase-quartz-biotite +/- chlorite +/- amphibole phyllite, schist, and gneiss representing metamorphosed volcanic tuffs and flows.

- Osp1** Quartz-plagioclase +/- biotite +/- muscovite granofels.

- Osp2** Dark gray amphibolite, locally containing garnet.

- Ooc** Cape Elizabeth Formation: Thin-bedded quartz-plagioclase-biotite-muscovite schist and granofels.

- Ooc1** Rusty weathering muscovite-biotite-garnet-staurolite schist and phyllite.

- Ooc2** Quartzose plagioclase-biotite phyllite with garnet-rich granofels (cotectic) beds. (On Western Horse Island, Casco Bay.)

- Ooc3** Cushing Formation: Medium-gray to light-gray, massive to thin-bedded, quartz-feldspar-biotite gneisses and subordinate schists representing metamorphosed pyroclastic volcanics and volcanogenic sediments. Includes varieties with garnet, hornblende, or relict blue quartz phenocryst fragments.

- Ooc4** Rusty weathering, fine-grained, plagioclase-quartz-muscovite schist.

- Ooc5** Wilson Cove Member: Very sulfidic, rusty weathering rocks, including garnet-biotite-amphibole gneiss and quartz-muscovite schist.

- Ooc6** Impure marble.

- Ooc7** Amphibolite with minor calc-silicate gneiss and impure marble.

- Ooc8** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc9** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc10** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc11** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc12** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc13** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc14** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc15** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc16** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

- Ooc17** Merpoint Member: Sulfidic quartz-plagioclase-muscovite-biotite schist.

Geologic editors:
Henry N. Berry IV and Arthur M. Hussey II

Contributing geologists:
Henry N. Berry IV, John W. Creasy, Timothy J. Cronan, Matthew R. Engelman, Richard A. Gilman, Chris Guzowski, Arthur M. Hussey II, Robert G. Marvinney, Alexander C. Robinson, Adam Schoonmaker, and Pauline M. Shyka

Cartographic design and production, by:
Robert D. Tucker, Bennett J. Wilson, Jr., and Robert A. Johnston

Robert G. Marvinney, State Geologist

Funding for the preparation of this map was provided in part by the U.S. Geological Survey National Geologic Mapping Program, Cooperative Agreement No. 1434-HQ-96-AG-01493.

Maine Geological Survey

Address: 22 State House Station, Augusta, Maine 04333
Telephone: 207-287-2801 E-mail: nrmm@state.me.us
Home page: <http://www.state.me.us/dnr/nrmm/nrmm.htm>

Open-File No. 98-1
1998

EXPLANATION OF SYMBOLS

Mesozoic Intrusions

- Basalt or diabase dike.

Isolated Paleozoic Intrusions

- Pegmatite.
- Aplitic.
- Muscovite-biotite granite.
- Garnet-bearing granite.
- Foliated biotite granite.

Isolated Paleozoic Metamorphic Rock Occurrences

- Schist.
- Calc-silicate rock.
- Quartz-feldspar-biotite granofels.

EXPLANATION OF PATTERNS

- Intensive rock mixed with abundant metamorphic rock.
- Syenite with porphyritic texture.

EXPLANATION OF LINES

- Approximate migmatite boundary. Separates moderately to strongly migmatitic rocks from weakly or non-migmatitic rocks. "m" toward the migmatitic side. Applies only to stratified rocks in the northeast part of the map. Migmatitic rocks to the west are not delineated separately.
- Intrusive or stratigraphic contact. Location inferred from nearest mapped bedrock exposures. Queried where not mapped.
- Fault, interpreted as a thrust fault. Offset inferred from deformed rock features, mainly brittle fabrics, or from apparently discordant structural or stratigraphic features. Probably of Paleozoic age.
- Fault, interpreted as a high-angle fault. Offset inferred from deformed rock features, mainly brittle fabrics, or from apparently discordant structural, metamorphic, or stratigraphic features. Of Paleozoic(?) to Mesozoic age.